

Interplay between Zeeman Coupling and Swap Action in Spin-based Quantum Computer Models: Error Correction in Inhomogeneous Magnetic Fields

Xuedong Hu, Rogerio de Sousa, and S. Das Sarma

Department of Physics, University of Maryland, College Park, MD 20742-4111

(February 1, 2008)

We consider theoretically the interplay between Zeeman coupling and exchange-induced swap action in spin-based quantum dot quantum computer models in the presence of inhomogeneous magnetic fields, which are invariably present in real systems. We estimate quantitatively swap errors caused by the inhomogeneous field, establishing that error correction would, in principle, be possible in the presence of non-uniform magnetic fields in realistic structures.

PACS numbers: 03.67.Lx

In recent years quantum computing has attracted widespread attention. The computers based on the principles of quantum mechanics, such as quantum parallelism and entanglement, promise to deliver results much faster than classical computers in certain tasks such as factoring and searching [1]. The impetus for constructing real quantum computer architectures arose from the seminal results [2] establishing that quantum error correction is theoretically possible and therefore decoherence is not an insurmountable barrier as was assumed earlier. There have been many proposals for quantum computer (QC) architectures based on various physical two-level systems, such as those using laser-cooled trapped ions [3,4], photons or atoms trapped in cavities [5,6], nuclear spins in bulk solutions [7] and crystal lattices [8], spins of electrons trapped in quantum dots [9,10] or donors [11], spins of donor nuclei in Si [12], superconducting devices [13], and others. The minimal requirements for a QC architecture are the existence of fundamental quantum bits (qubit) and the ability to carry out single- and two-qubit operations such as swap and controlled-NOT, as well as suitable hardwares for reading (input) and writing (output or measurement) operations. Among the many obstacles facing the successful demonstration of nontrivial quantum computation in specific QC hardwares, the most daunting are the problem of quantum decoherence and the difficulty in achieving precise control over the various unitary operations necessary for quantum computation. In this Letter we consider the theoretical issue of controlling the swap operation in the proposed solid state QC architecture involving quantum dot spin entanglement, with externally-controlled electrostatic gates and magnetic fields providing the unitary operation.

Initial work on QC hardwares concentrated on atomic/molecular systems partly because well-defined single-qubit operations are comparatively easier to control in atomic systems such as trapped ions [3,4]. Successful QC architectures will eventually require many qubits working in parallel, which would be difficult to achieve in atomic systems. Proposed solid state QC architectures

are, in principle, scalable to many qubits, although no one has yet successfully demonstrated a controlled single-qubit operation in solid state QC systems. This is a curious dichotomy in the current QC hardware research—the architectures with demonstrated single-qubit operations are difficult to scale up while the presumably scalable solid state architectures have not yet been able to demonstrate single-qubit control. It is thus crucial to explore the challenges facing coherent control of qubits in solid state structures, particularly the issue of possible error corrections in realistic systems. Among various microscopic degrees of freedom that have been considered for the role of qubits in solid state QC architectures, spins of electrons or nuclei are natural candidates because of their well-defined Hilbert spaces and their relatively long decoherence time compared to the orbital degrees of freedom. In the proposed spin-based QC architectures, exchange interaction plays a fundamental role of establishing two-qubit entanglement [8,9,11,12], while Zeeman coupling to an external magnetic field provides various single-qubit operations. Generally Zeeman coupling is treated separately from the exchange interaction [14,15] because the latter originates from the Coulomb interaction (due to Pauli principle) while the former is purely a spin effect, and the two interaction terms in the Hamiltonian commute with each other if the Zeeman term is homogeneous. However, solid state heterostructures are intrinsically inhomogeneous and magnetic imperfections or impurities are likely to be present, leading to inhomogeneous stray magnetic fields. Furthermore, parallel pulse schemes, in which exchange interaction and inhomogeneous Zeeman coupling are present simultaneously, have been proposed to expedite operations of spin-based QCs [16]. Therefore, it is necessary to explore the interplay between exchange interaction and Zeeman coupling in relation to spin-based QC architectures, as we do here.

In this Letter we focus on the effects of Zeeman coupling when the external magnetic field is spatially inhomogeneous. Zeeman coupling is generally neglected in these studies [14,15] based partly on the assumption that

the effects are small and do not lead to any qualitative changes. While it is true in uniform fields, this assumption may not hold in an inhomogeneous magnetic field. The key motivations for our study are the observations that spatially inhomogeneous fields are intrinsic to spin-based solid state QC architectures and that for parallel schemes the magnetic fields at the locations of the two spins are generally different [16], which means the total field must be inhomogeneous. If only the single-spin evolution is involved, effects of local magnetic field can be corrected in such single-qubit operations by techniques such as spin echoes [17]. However, if two-spin entangled evolution is also involved, it is not clear whether errors caused by inhomogeneous fields can still be eliminated, *i.e.* whether such errors lie within the current QC error correction constraints.

As an example we study how Zeeman coupling may affect the proposed operations of spin-based quantum dot quantum computers (QDQC). Our results, however, are quite general and can be applied to other spin-based models with finite magnetic fields. In particular, we study the effect of a finite inhomogeneous magnetic field on swap actions. For most of the spin-based schemes, swap action U_{sw} , in which two spins exchange their states, is one of the most basic operations. It is used to construct conditional phase shifts (CPS), $U_{CPS} = e^{i\frac{\pi}{4}\sigma_{1z}} e^{-i\frac{\pi}{4}\sigma_{2z}} U_{sw}^{\frac{1}{2}} e^{i\frac{\pi}{4}\sigma_{1z}} U_{sw}^{\frac{1}{2}}$ (with σ being the Pauli matrices for spins), which can then be converted to controlled-NOT (CNOT) easily [9]. In addition, swaps are used to move spin states around [9,12] so that an arbitrary pair of spins can be brought into controlled entanglement, which is essential to quantum computation [1]. It is therefore important to investigate how Zeeman splitting will affect the swap gate and what are the consequences if these effects are non-trivial.

In a single envelope function approach [15], the Hamiltonian for two electrons trapped in a lateral double-quantum-dot is

$$H = \sum_i^2 \left[\frac{1}{2m^*} \left(\mathbf{p} + \frac{e}{c} \mathbf{A}(\mathbf{r}_i) \right)^2 + V(\mathbf{r}_i) + g^* \mu_B \mathbf{B}(\mathbf{r}_i) \cdot \mathbf{S}_i \right] + \frac{e^2}{\epsilon r_{12}}. \quad (1)$$

Notice that here the magnetic field has a spatial dependence. The Hamiltonian can be expanded in a basis of two-spin eigenstates (spin singlet and triplet states). If the magnetic field is uniform, the Zeeman coupling depends only on the total spin along the field direction and commutes with the total spin operator $\mathbf{S}^2$, so that singlet and triplet states remain eigenstates of the two-electron system. On the other hand, if the magnetic field is inhomogeneous, it destroys the symmetry between the two spins, and the singlet state and one of the triplet states (the one with $S_n = 0$, where $\mathbf{n}$

is the field direction) couple. For example the coupling between the lowest singlet and triplet states (in the Heitler-London approximation) is $\langle T | V_Z | S \rangle = g\mu_B/2 \times \{ \langle L | B_z(\mathbf{r}) | L \rangle - \langle R | B_z(\mathbf{r}) | R \rangle + \langle L | B_z(\mathbf{r}) | R \rangle \langle R | L \rangle - \langle R | B_z(\mathbf{r}) | L \rangle \langle L | R \rangle \}$, where the singlet state is $|S\rangle = [|L(1)\rangle |R(2)\rangle + |L(2)\rangle |R(1)\rangle] (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)/2$, the triplet state is $|T\rangle = [|L(1)\rangle |R(2)\rangle - |L(2)\rangle |R(1)\rangle] (|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)/2$, and the Zeeman coupling is $V_Z = g\mu_B [B_z(\mathbf{r}_1)S_{1z} + B_z(\mathbf{r}_2)S_{2z}]$. Here we have assumed that the external field is along the z direction. $|L\rangle$ and $|R\rangle$ are the left and right quantum dot ground states. Therefore, the inhomogeneous magnetic field couples the singlet and the $S_z = 0$ triplet states so that two of the eigenstates of the system are no longer the eigenstates of the total spin. When the overlap between the two ground state wavefunctions is small, the main contribution to this coupling comes from the average field difference between the two quantum dots. Below we explore the consequences of this loss of symmetry.

When the overlap of the electronic wavefunctions is small, we can assume that the electron orbital degrees of freedom are frozen, so that an effective spin Hamiltonian quite faithfully describes the two-spin system [14,15]:

$$H_s = J \mathbf{S}_1 \cdot \mathbf{S}_2 + \gamma_1 S_{1z} + \gamma_2 S_{2z}, \quad (2)$$

where $\gamma_1 = g\mu_B \langle L | B_z(\mathbf{r}) | L \rangle$ and $\gamma_2 = g\mu_B \langle R | B_z(\mathbf{r}) | R \rangle$ are local Zeeman couplings due to applied or stray magnetic fields. Here we have implicitly assumed, based on the small interdot wavefunction overlap, that the two spins are distinguishable, with spin 1 on the left dot and spin 2 on the right dot. We have also assumed that the field is entirely along the z direction. Whether such a choice is reasonable will be discussed later.

Hamiltonian (2) can be expressed in the basis of four two-spin states $|\uparrow\uparrow\rangle$, $|\uparrow\downarrow\rangle$, $|\downarrow\uparrow\rangle$, and $|\downarrow\downarrow\rangle$, and its eigenstates can be easily obtained. The two polarized states are decoupled from the other two, which are mixtures of singlet and $S_z = 0$ triplet states: $|\psi_1\rangle = |\uparrow\uparrow\rangle$, $|\psi_2\rangle = |\downarrow\downarrow\rangle$, $|\psi_3\rangle = c_1 |\uparrow\downarrow\rangle + c_2 |\downarrow\uparrow\rangle$, and $|\psi_4\rangle = c_2 |\uparrow\downarrow\rangle - c_1 |\downarrow\uparrow\rangle$. Here the coefficients c_1 and c_2 satisfy the relations $c_2/c_1 = \sqrt{1 + (\delta/2J)^2} - \delta/2J$ and $c_1^2 + c_2^2 = 1$, where $\delta = \gamma_1 - \gamma_2$ represents the field inhomogeneity. The energies of the latter two states are also shifted from those of singlet and triplet states (with $\tilde{\delta}$ being the shift): $E_1 = J + \Delta$, $E_2 = J - \Delta$, $E_3 = -J + \sqrt{4J^2 + \delta^2} = J + \tilde{\delta}$, and $E_4 = -J - \sqrt{4J^2 + \delta^2} = -J - \tilde{\delta}$, where $\Delta = \gamma_1 + \gamma_2$ represents the average magnetic field. An important question here is whether these mixtures and shifts will cause any error in quantum computation in the schemes based on the exchange interaction. After all, the swap action in these models depends on the perfect phase matching in the evolution of singlet and triplet states, as we will show below. Since swap operation is an essential component of a spin-based QDQC and several other architectures, we need to precisely quantify the effects of mixtures in singlet and triplet states on the swap action.

To determine whether swap is affected, we explore whether a product state of spins 1 and 2 will evolve into a product state (pure states for both spins) again. Our strategy here is to calculate the Schmidt number of the two-spin state. We will evolve our two-spin state, calculate its density matrix, trace out the second spin so that the density matrix of the first spin is left. We can then look for the eigenvalues of this density matrix. If at some time it has only one non-vanishing eigenvalue, the state of the first spin is pure. We can then find out whether this pure state corresponds to a swapped state.

Our initial state is a product state given by

$$|\phi(0)\rangle = (\alpha_1|\uparrow\rangle + \alpha_2|\downarrow\rangle)(\beta_1|\uparrow\rangle + \beta_2|\downarrow\rangle). \quad (3)$$

If the two electrons are located in two well-separated quantum dots in the beginning, the above product state does not violate the antisymmetry requirement of a two-fermion state. This state can be expanded in the basis of the eigenstates of Hamiltonian (2). It then evolves under Hamiltonian (2). The two-spin state at time t takes the form $|\phi(t)\rangle = \alpha_1\beta_1 e^{-iE_1t/\hbar}|\uparrow\uparrow\rangle + \alpha_2\beta_2 e^{-iE_2t/\hbar}|\downarrow\downarrow\rangle + (\alpha_1\beta_2c_1 + \alpha_2\beta_1c_2)e^{-iE_3t/\hbar}|\uparrow\downarrow\rangle + (\alpha_1\beta_2c_2 - \alpha_2\beta_1c_1)e^{-iE_4t/\hbar}|\downarrow\uparrow\rangle$. The corresponding density matrix for the first spin can then be calculated straightforwardly: $\rho_1 = \text{Tr}_2\{\rho_{12}\} = \text{Tr}_2\{|\phi(t)\rangle\langle\phi(t)|\}$. The matrix elements of ρ_1 are all functions of the constants $\alpha_1, \alpha_2, \beta_1, \beta_2, c_1, c_2$, field inhomogeneity δ , exchange coupling J , and time t ($\rho_{1\uparrow\downarrow}$ is also a function of the average field Δ). The eigenvalue equation for ρ_1 is

$$\lambda^2 - (\rho_{1\uparrow\uparrow} + \rho_{1\downarrow\downarrow})\lambda + (\rho_{1\uparrow\uparrow}\rho_{1\downarrow\downarrow} - |\rho_{1\uparrow\downarrow}|^2) = 0. \quad (4)$$

To have a pure state for the first spin, which means that only one eigenvalue of the density matrix ρ_1 is non-vanishing, the last term on the left-hand-side of the above equation has to vanish:

$$\rho_{1\uparrow\uparrow}\rho_{1\downarrow\downarrow} - |\rho_{1\uparrow\downarrow}|^2 = 0. \quad (5)$$

The explicit expression for Eq. (5) is quite complicated, so let us first look at the simple uniform field situation, when $\delta = 0$ and $c_1 = c_2 = 1/\sqrt{2}$. Equation (5) then becomes $|\alpha_1\beta_1\alpha_2\beta_2 - \frac{1}{4}[(\alpha_1\beta_2 + \alpha_2\beta_1) + (\alpha_1\beta_2 - \alpha_2\beta_1)e^{i\theta}][(\alpha_1\beta_2 + \alpha_2\beta_1) - (\alpha_1\beta_2 - \alpha_2\beta_1)e^{i\theta}]|^2 = 0$, where $\theta = 4Jt$. The solutions here are $e^{i\theta} = \pm 1$. When $e^{i\theta} = 1$, $|\phi_1(t)\rangle = \alpha_1|\uparrow\rangle + \alpha_2e^{i\Delta t}|\downarrow\rangle$, the state of the first spin returns to its initial state with a phase shift between the two coefficients. When $e^{i\theta} = -1$, $|\phi_1(t)\rangle = \beta_1e^{i\Delta t}|\uparrow\rangle + \beta_2|\downarrow\rangle$, the swap is achieved with the exception of an additional phase that can be corrected easily with a single-spin operation. For example, a pulse sequence can be constructed based on the phase-shifted swap U_{psw} to produce a CNOT gate

$$U_{CNOT} = e^{i\frac{\pi}{4}\sigma_{2y}} e^{i(\frac{\pi}{4} + \frac{\delta t}{2})\sigma_{1z}} e^{-i(\frac{\pi}{4} - \frac{\delta t}{2})\sigma_{2z}} \times U_{psw}^{\frac{1}{2}} e^{i\frac{\pi}{2}\sigma_{1z}} U_{psw}^{\frac{1}{2}} e^{-i\frac{\pi}{4}\sigma_{2y}}. \quad (6)$$

Physically, a uniform field means that the Zeeman coupling couples to the total spin (including both electron spins), so that the Zeeman term commutes with the exchange term in the Hamiltonian (2), and therefore does not change the eigenstates. The shifts in the energy levels of the polarized states cause additional phase shift, but can be corrected by applying an opposite magnetic field with the same pulse shape, magnitude, and length. In summary, an external uniform magnetic field does not qualitatively change the proposed QC algorithm. Logistically it makes the QC operation more difficult because of the necessary correction pulses.

If the magnetic field is inhomogeneous, $\delta \neq 0$. Let us look at a simple situation when the initial state is $|\phi(0)\rangle = |\uparrow\downarrow\rangle$, *i.e.* $\alpha_1 = \beta_2 = 1$ and $\alpha_2 = \beta_1 = 0$. Eq. (5) now takes the simple form of

$$|c_1c_2(1 - e^{i\theta})(c_1^2 + c_2^2e^{i\theta})|^2 = 0, \quad (7)$$

where $\theta = (4J + 2\delta)t$. When $c_1 \neq c_2$, as is the case when $\delta \neq 0$, the only solution of this equation is $e^{i\theta} = 1$, which corresponds to the “return” operation with an additional phase: $|\phi_1(t)\rangle = \alpha_1|\uparrow\rangle + \alpha_2e^{i\Delta t}|\downarrow\rangle$. The condition for an exact swap does not exist anymore. To find the best approximation to a swap (so that the state of first spin is as close as possible to being “down” in the current special case), we calculate the minima of the expression in Eq. (7) and find that $e^{i\theta} = -1$ does still produce a minimum in $|(1 - e^{i\theta})(c_1^2 + c_2^2e^{i\theta})|^2$ (which is, however, no longer zero). When this condition is satisfied, the density matrix of the first spin is

$$\rho_1|_{e^{i\theta}=-1} = \frac{1}{1+x^2}|\downarrow\rangle\langle\downarrow| + \frac{x^2}{1+x^2}|\uparrow\rangle\langle\uparrow|, \quad (8)$$

where $x = \delta/(2J)$. Therefore, the state of the first spin cannot be simultaneously pure and the same as the initial state of the second spin. The state of the first spin will remain mixed (when it is close to a swap) with the state of the second spin. According to Eq. (8), there would be an error of the magnitude x^2 if we perform such a “swap” operation, which needs to be corrected.

For GaAs-based QDQC, the Zeeman splitting is $g\mu_B \approx 2.55 \times 10^{-2}$ meV/Tesla, while a single Bohr magneton produces a field of about 5 Gauss at 1 nm distance and almost nothing at a distance of 100 nm in GaAs. This difference in the magnetic field leads to a x^2 in the order of 10^{-6} , which is right within the capability of currently available error correction schemes. Since the field involved in δ is the average value over an entire quantum dot, local magnetic field inhomogeneity caused by impurities should not cause intractable errors.

Mathematically, inhomogeneity in the magnetic field means that a part of the Hamiltonian (2) does not commute with the exchange term $JS_1 \cdot S_2$, thus the eigenstates will change, leading to errors in swap, which will cause errors in CNOT in the proposed sequential scheme

[9]. In the parallel pulse scheme [16], conditional phase shifts can be produced directly from the Hamiltonian (2) (with precisely controlled field inhomogeneity), circumventing the swap actions. However, stray fields, which may be invariably present in realistic solid state QDQC architectures due to magnetic impurities/imperfections, can still produce errors which need to be corrected (even in solely exchange-based models which have no applied magnetic field [18]). Furthermore, the impossibility of exact swap does make the transfer of spins through swap an error-prone process in the presence of an inhomogeneous field.

Finally, inhomogeneity in the external magnetic field is not only a possible error source in swap and other two-qubit operations, but also an important factor in the single-qubit operations determined solely by the Zeeman coupling. Here the main concern is the precise definition of the direction of the quantization axis for the electron spins. For example, if spin-up and -down along z direction correspond to the two states of our qubit, then the z axis is our quantization axis which provides a reference frame for all the single-qubit operations. In our calculations the magnetic field is purely along the z direction. The only spatial variation is in its magnitude. However, according to Maxwell equations, in a steady state, $\nabla \times \mathbf{B} = 0$. Thus, if $\partial B_z / \partial x \neq 0$, $\partial B_x / \partial z = \partial B_z / \partial x \neq 0$. If we have a perfect two dimensional QD system lying entirely in the x - y plane, we can assume that B_x happens to vanish in the plane of the quantum dots. However, using the example of GaAs quantum well QD's, the thickness of the well is generally around 5 – 10 nm, which characterizes the typical z -width of the QD's. Therefore, to have a vanishing Zeeman coupling along x direction, the B_x field has to vanish along the middle plane of the quantum well and be an odd function in the z (growth) direction, which is a quite stringent constraint. If we choose z -direction as our quantizing axis, a finite B_x field, which may be unavoidable, would tend to flip the spins. For a 1 Gauss B_x field, this flipping rate is about 0.65 MHz, corresponding to a flipping time of 1.5 μ s. Considering that the gates should be operated as slowly as possible to satisfy the adiabatic condition, this is a constraint one has to take into consideration. In fact, “decoherence” induced by a fluctuating inhomogeneous transverse field may turn out to be an important extrinsic decoherence channel for QDQC operations.

This work is supported by ARDA, LPS, and US-ONR.

- and D.P. DiVincenzo, *Nature* **404**, 247 (2000).
- [2] P. Shor, *Phys. Rev. A* **52**, R2493 (1995); A. Steane, *Phys. Rev. Lett.* **77**, 793 (1996).
 - [3] J.I. Cirac and P. Zoller, *Phys. Rev. Lett.* **74**, 4091 (1995).
 - [4] C. Monroe et al., *Phys. Rev. Lett.* **75**, 4714 (1995).
 - [5] T. Sleator and H. Weinfurter, *Phys. Rev. Lett.* **74**, 4087 (1995).
 - [6] Q.A. Turchette et al., *Phys. Rev. Lett.* **75**, 4710 (1995).
 - [7] D.G. Cory, A.F. Fahmy, and T.F. Havel, *Proc. Natl. Acad. Sci. U.S.A.* **94**, 1634 (1997); N.A. Gershenfeld and I.L. Chuang, *Science* **275**, 350 (1997).
 - [8] F. Yamaguchi and Y. Yamamoto, *Appl. Phys. A* **68**, 1 (1999).
 - [9] D. Loss and D.P. DiVincenzo, *Phys. Rev. A* **57**, 120 (1998).
 - [10] A. Imamoglu et al., *Phys. Rev. Lett.* **83**, 4204 (1999).
 - [11] R. Vrijen et al., *Phys. Rev. A* **62**, 012306 (2000).
 - [12] B.E. Kane, *Nature (London)* **393**, 133 (1998).
 - [13] A. Shnirman, G. Schön, and Z. Hermon, *Phys. Rev. Lett.* **79**, 2371 (1997).
 - [14] G. Burkard, D. Loss, and D.P. DiVincenzo, *Phys. Rev. B* **59**, 2070 (1999).
 - [15] X. Hu and S. Das Sarma, *Phys. Rev. A* **61**, 062301 (2000).
 - [16] G. Burkard et al., *Phys. Rev. B* **60**, 11404 (1999).
 - [17] C.P. Slichter, *Principles of Magnetic Resonance* (Springer-Verlag, Berlin, 1989).
 - [18] D.P. DiVincenzo, D. Bacon, J. Kempe, G. Burkard, and K.B. Whaley, *Nature* **408**, 339 (2000).

[1] A. Ekert and R. Jozsa, *Rev. Mod. Phys.* **68**, 733 (1996); A. Steane, *Rep. Prog. Phys.* **61**, 117 (1998); C.H. Bennett